

QCD sum rules and thermal properties of Charmonium in the vector channel

C. A. Dominguez^{a,b,1}, M. Loewe^{c,2,3,*}, J. C. Rojas^{d,2}, Y. Zhang^b

^aCentre for Theoretical Physics and Astrophysics, University of Cape Town, Rondebosch 7700, South Africa

^bDepartment of Physics, Stellenbosch University, Stellenbosch 7600, South Africa.

^cFacultad de Física, Pontificia Universidad Católica de Chile, Casilla 306, Santiago 22, Chile.

^dDepartamento de Física, Universidad Católica del Norte, Casilla 1280, Antofagasta, Chile.

Abstract

The thermal evolution of the hadronic parameters of charmonium in the vector channel, i.e. the J/ψ resonance mass, coupling (leptonic decay constant), total width, and continuum threshold is analyzed in the framework of thermal Hilbert moment QCD sum rules. The continuum threshold s_0 , as in other hadronic channels, decreases with increasing temperature until the PQCD threshold $s_0 = 4m_Q^2$ is reached at $T \simeq 1.22 T_c$ (m_Q is the charm quark mass) and the J/ψ mass is essentially constant in a wide range of temperatures. The other hadronic parameters behave in a very different way from those of light-light and heavy-light quark systems. The total width grows with temperature up to $T \simeq 1.04 T_c$ beyond which it decreases sharply with increasing T . The resonance coupling is also initially constant beginning to increase monotonically around $T \simeq T_c$. This behavior strongly suggests that the J/ψ resonance might survive beyond the critical temperature for deconfinement, in agreement with lattice QCD results.

Keywords: Finite temperature field theory, hadron physics.

1. Introduction

We discuss here the thermal evolution of the hadronic parameters of J/ψ in the vector channel, using thermal QCD Sum Rules [1]. We refer the reader to the original article [2] for details. This technique has been used previously in the light-light and in the heavy-light quark sector [3]- [5], with the following emerging picture: (i) For increasing temperature, hadronically stable particles develop a non-zero width, and resonances become broader, diverging at a critical temperature interpreted as the deconfinement temperature (T_c). The thermal resonance broadening was first proposed in [6]. ii) Above the resonance region, the continuum threshold in hadronic spectral functions, i.e. the onset of perturbative QCD (PQCD), decreases monotonically with increasing temperature. When $T \rightarrow T_c$ hadrons disappear from the spectrum. (iii) This scenario is also supported

by the behavior of hadronic couplings, or leptonic decay constants, which approach zero as $T \rightarrow T_c$. Masses, on the other hand, do not provide information about deconfinement.

The thermal behavior of the heavy-heavy quark correlator should be different from that involving at least one light quark since: a) In the light-light and heavy-light quark sector, the PQCD contribution is dominated by the time-like spectral function (annihilation term), which is relatively unimportant in relation to the light quark condensate contribution, being the scattering PQCD spectral function highly suppressed. Instead, for heavy-heavy quark systems this term becomes increasingly important with increasing temperature while the annihilation term only contributes near threshold; b) The non-perturbative QCD sector in the operator product expansion (OPE) of light-light and heavy-light quark correlators is driven by the light quark condensate, responsible for the behavior of the continuum threshold since $s_0(T)/s_0(0) \simeq \langle \bar{q}q \rangle / \langle \bar{q}q \rangle$ [4]-[7]. The light quark condensate is the order parameter for chiral symmetry restoration. In contrast, for heavy-heavy quark correlators the leading power correction in the OPE is that of the gluon condensate, which has a very different thermal behavior. In this approach, the criti-

*Speaker

Email addresses: Cesareo.Dominguez@uct.ac.za (C. A. Dominguez), mloewe@fis.puc.cl (M. Loewe), jurojas@ucn.cl (J. C. Rojas), Yingwen.Zhang@uct.ac.za (Y. Zhang)

¹Supported by NRF South Africa

²Supported by Fondecyt Chile under grant 1095217

³Supported by Proyecto Anillos ACT119

cal temperature for deconfinement is a phenomenological parameter which does not need to coincide with e.g. the critical temperature obtained in lattice QCD [8]. In fact, results from QCD sum rules lead to values of T_c somewhat lower than those from lattice QCD. In order to compare with other approaches, we express our results in terms of the ratio T/T_c .

We find for charmonium in the vector channel that the continuum threshold, $s_0(T)$, decreases with increasing T , being driven by the gluon condensate and the PQCD spectral function in the space-like region, until it reaches the PQCD threshold $s_0 = 4m_Q^2$ at $T \simeq 1.22 T_c$ (m_Q is the charm quark mass). Below this value of s_0 the sum rules cease to be valid. The J/ψ mass remains basically constant as in the light-light or heavy-light systems. We have, however, a very different thermal evolution of the width and the coupling. Both are almost independent of T up to $T \simeq 0.8 T_c$ where the width begins to increase substantially, but then above $T \simeq 1.04 T_c$ it starts to decrease sharply, and the coupling increases also sharply. This suggests the survival of the J/ψ resonance above the deconfinement temperature.

The PQCD spectral function in the space-like region plays here a very important role. Non-relativistic approaches to charmonium at finite temperature would normally miss this contribution. In fact, the complex energy plane in the non-relativistic case would only have one cut along the positive real axis, which would correspond to the time-like (annihilation) region of PQCD. The space-like contribution ($q^2 = (\omega^2 - |\mathbf{q}|^2) \leq 0$) in the form of a cut in the energy plane centered at the origin for $-|\mathbf{q}| \leq \omega \leq |\mathbf{q}|$, would not be present in the non-relativistic case.

2. Hilbert Moment QCD Sum Rules

We consider the correlator of the heavy-heavy quark vector current at finite temperature

$$\begin{aligned} \Pi_{\mu\nu}(q^2, T) &= -(g_{\mu\nu}q^2 - q_\mu q_\nu) \Pi(q^2, T) \\ &= i \int d^4x e^{iqx} \theta(x_0) \langle [V_\mu(x), V_\nu^\dagger(0)] \rangle, \end{aligned} \quad (1)$$

where $V_\mu(x) =: \bar{Q}(x)\gamma_\mu Q(x)$:, and $Q(x)$ is the heavy (charm) quark field. The matrix element above is understood to be the Gibbs average in the quark-gluon basis. The imaginary part of the vector correlator in PQCD at finite temperature involves two pieces, one in the time-like region ($q^2 \geq 4m_Q^2$), $\text{Im} \Pi_a(q^2, T)$, which survives at $T=0$, and one in the space-like region ($q^2 \leq 0$), $\text{Im} \Pi_s(q^2, T)$, which vanishes at $T=0$. To leading order in PQCD we find

$$\begin{aligned} \frac{1}{\pi} \text{Im} \Pi_a(q^2, T) &= \frac{3}{16\pi^2} \int_{-\nu}^{\nu} dx (1-x^2) \\ &\quad \left[1 - n_F\left(\frac{|\mathbf{q}|x + \omega}{2T}\right) - n_F\left(\frac{|\mathbf{q}|x - \omega}{2T}\right) \right], \end{aligned} \quad (2)$$

where $\nu^2 = 1 - 4m_Q^2/q^2$, m_Q is the heavy quark mass, $q^2 = \omega^2 - \mathbf{q}^2 \geq 4m_Q^2$, and $n_F(z) = (1 + e^z)^{-1}$ is the Fermi thermal function. In the rest frame of the thermal bath, $|\mathbf{q}| \rightarrow 0$, the above result reduces to

$$\begin{aligned} \frac{1}{\pi} \text{Im} \Pi_a(\omega, T) &= \frac{\nu(3-\nu^2)}{8\pi^2} [1 - 2n_F(\omega/2T)] \\ &\quad \times \theta(\omega - 2m_Q). \end{aligned} \quad (3)$$

The quark mass is assumed independent of T , a good approximation for $T < 200$ MeV [9]. Only the leading order in the strong coupling will be considered here.

The PQCD piece in the space-like region demands a careful analysis. In the complex energy plane, and in the space-like region, the correlator $\Pi(q^2)$, Eq.(1), has a cut centered at the origin and extending between $\omega = -|\mathbf{q}|$ and $\omega = |\mathbf{q}|$. In the rest frame this cut produces a delta function $\delta(\omega^2)$ in the imaginary part of $\Pi(q^2)$. The result is

$$\begin{aligned} \frac{1}{\pi} \text{Im} \Pi_s(\omega, T) &= \frac{2}{\pi^2} m_Q^2 \delta(\omega^2) \times \\ &\quad \left[n_F\left(\frac{m_Q}{T}\right) + \frac{2T^2}{m_Q^2} \int_{m_Q/T}^{\infty} y n_F(y) dy \right]. \end{aligned} \quad (4)$$

The corresponding hadronic representation is parametrized in terms of the ground state resonance, the J/ψ , followed by a continuum given by PQCD after a threshold $s_0 > M_V^2$. In the zero width approximation, the hadronic spectral function is

$$\begin{aligned} \frac{1}{\pi} \text{Im} \Pi(s, T)|_{HAD} &= \frac{1}{\pi} \text{Im} \Pi(s, T)|_{RES} \theta(s_0 - s) \\ &\quad + \frac{1}{\pi} \text{Im} \Pi(s, T)|_{PQCD} \theta(s - s_0) \\ &= 2 f_V^2(T) \delta(s - M_V^2(T)) + \frac{1}{\pi} \text{Im} \Pi(s, T)_a \theta(s - s_0), \end{aligned} \quad (5)$$

where $s \equiv q^2 = \omega^2 - \mathbf{q}^2$. The leptonic decay constant is defined as $\langle 0 | V_\mu(0) | V(k) \rangle = \sqrt{2} M_V f_V \epsilon_\mu$.

When considering a finite (total) width the following replacement will be understood

$$\pi \delta(s - M_V^2(T)) \rightarrow \frac{M_V(T) \Gamma_V(T)}{(s - M_V^2(T))^2 + M_V^2(T) \Gamma_V^2(T)}, \quad (6)$$

The hadronic scattering term, due to current scattering off D-mesons, is negligible [2]. The correlation function $\Pi(q^2, T)$, Eq.(1), satisfies a once subtracted dispersion relation. To eliminate the subtraction one can use Hilbert moments, i.e.

$$\begin{aligned}\varphi_N(Q^2, T) &\equiv \frac{(-)^N}{(N)!} \left(\frac{d}{dQ^2}\right)^N \Pi(Q^2, T) \\ &= \frac{1}{\pi} \int_0^\infty \frac{ds}{(s + Q^2)^{N+1}} \text{Im} \Pi(s, T),\end{aligned}\quad (7)$$

where $N = 1, 2, \dots$, and $Q^2 \geq 0$ is an external four-momentum squared, to be considered as a free parameter. Using Cauchy's theorem in the complex s -plane, the Hilbert moments become Finite Energy QCD sum rules (FESR), i.e.

$$\varphi_N(Q^2, T)|_{RES} = \varphi_N(Q^2, T)|_{QCD}, \quad (8)$$

where

$$\varphi_N(Q^2, T)|_{RES} \equiv \frac{1}{\pi} \int_0^{s_0(T)} \frac{ds}{(s + Q^2)^{N+1}} \text{Im} \Pi(s, T)|_{RES}, \quad (9)$$

$$\begin{aligned}\varphi_N(Q^2, T)|_{QCD} &\equiv \frac{1}{\pi} \int_{4m_Q^2}^{s_0(T)} \frac{ds}{(s + Q^2)^{N+1}} \text{Im} \Pi_a(s, T) \\ &+ \frac{1}{\pi} \int_0^\infty \frac{ds}{(s + Q^2)^{N+1}} \text{Im} \Pi_s(s, T) + \varphi_N(Q^2, T)|_{NP},\end{aligned}\quad (10)$$

and $\text{Im} \Pi(s, T)|_{RES}$ is given by the first term in Eq.(5) modified in finite-width according to Eq.(6), and the PQCD spectral functions are given by Eqs.(3) and (4).

The dimension $d=4$ non perturbative term in the OPE is well known in the literature, see [2] for details. The dependence on N is quite cumbersome and it is proportional to the gluon condensate $\langle\langle \frac{\alpha_s}{\pi} G^2 \rangle\rangle$. At low temperatures, this condensate

has been calculated in chiral perturbation theory [10]. In this framework the condensate remains essentially constant up to $T \sim T_c \simeq 100$ MeV, after which it decreases sharply. In order to go beyond the low temperature regime of chiral perturbation theory, lattice QCD provides the right tool. A good approximation [11] is given by the expression

$$\langle\langle \frac{\alpha_s}{\pi} G^2 \rangle\rangle = \left\langle \frac{\alpha_s}{\pi} G^2 \right\rangle \left[\theta(T^* - T) + \frac{1 - \frac{T}{T_c^*}}{1 - \frac{T^*}{T_c^*}} \theta(T - T^*) \right] \quad (11)$$

where $T^* \approx 150$ MeV is the breakpoint temperature where the condensate begins to decrease appreciably, and $T_c^* \approx 250$ MeV is the temperature at which $\langle\langle \frac{\alpha_s}{\pi} G^2 \rangle\rangle_{T_c} = 0$.

Returning to the Q^2 dependence of the Hilbert moments, Eq.(7), we shall fix Q^2 and $s_0(0)$ from the experimental values of the mass, the coupling, and the width at $T=0$. At finite temperature there are non-diagonal (Lorentz non-invariant) condensates that might contribute to the OPE. Non-gluonic operators are highly suppressed [5], [12] so that they can be safely ignored. We have considered also a gluonic twist-two term in the OPE introduced in [13], and computed on the lattice in [14]. Its impact is small, (2-6)%, and plays no appreciable role in the results.

3. Results

We begin by determining s_0 and Q^2 at $T=0$ from the moments, Eq.(8), and using as input the experimental values [15] $M_V = 3.097$ GeV, $f_V = 196$ MeV, and $\Gamma_V = 93.2$ keV, as well as $m_Q = 1.3$ GeV, and [16] $\langle 0 | \frac{\alpha_s}{12\pi} G^2 | 0 \rangle \simeq 5 \times 10^{-3} \text{ GeV}^4$. In the zero-width approximation one finds from Eq.(9) that

$$\frac{\varphi_1(Q^2)|_{RES}}{\varphi_2(Q^2)|_{RES}} = \frac{\varphi_2(Q^2)|_{RES}}{\varphi_3(Q^2)|_{RES}}. \quad (12)$$

Given the extremely small total width of the J/ψ it turns out that the above relation also holds with extreme accuracy in finite width. Using Eq.(8) this leads to

$$\frac{\varphi_1(Q^2)|_{QCD}}{\varphi_2(Q^2)|_{QCD}} = \frac{\varphi_2(Q^2)|_{QCD}}{\varphi_3(Q^2)|_{QCD}}, \quad (13)$$

which depends only on the two unknowns s_0 and Q^2 , and provides the first equation to determine this pair of parameters. The second equation can be e.g. Eq.(8) with $N = 1$. In this way we find that $s_0 = 11.64 \text{ GeV}^2$, and $Q^2 = 10 \text{ GeV}^2$ reproduce the experimental values of the mass, coupling, and width of J/ψ within less than 1%. This whole set of hadronic parameters will then be used to normalize the corresponding parameters at finite temperature. In this way, see [2] for details, we were able to find the thermal evolution of s_0 , the J/ψ mass, its width and its coupling. We show here only the behavior of the width and the coupling (Figs. 1 and 2) since these are the most important results of this analysis.

Both the width and the coupling can only be determined up to $T_f \simeq 1.1 T_c$ beyond which $s_0(T) < M_V^2(T)$ and the FESR integrals have no longer a support. The temperature behavior of the width and the coupling shown in Figs. 1 and 2 strongly suggests the survival




Figure 1: The ratio $\Gamma_V(T)/\Gamma_V(0)$ as a function of T/T_c .




Figure 2: The ratio $f_V(T)/f_V(0)$ as a function of T/T_c .

of the J/ψ above the critical temperature for deconfinement. This conclusion agrees with results from lattice QCD [8], but disagrees with non-relativistic determinations. As pointed out earlier, the reason for this disagreement might very well be the absence of the central cut (QCD scattering term) in the energy plane in non-relativistic frameworks.

References

- [1] A.I. Bochkarev and M.E. Shaposnikov, Nucl. Phys. B 286 (1986) 220.
- [2] C. A. Dominguez, M. Loewe, J. C. Rojas and Y. Zhang, Phys. Rev. D 81 (2010) 014007.
- [3] R.J. Furnstahl, T. Hatsuda and S.H. Lee, Phys. Rev. D 42 (1990) 1744; C. Adami, T. Hatsuda and I. Zahed, Phys. Rev. D 43(1991) 921; C. Adami and I. Zahed, Phys. Rev. D 45 (1992) 4312; T. Hatsuda, Y. Koike and S.-H. Lee, Phys. Rev. D 47 (1993) 1225; *ibid.* Nucl. Phys. B 394 (1993) 221; Y. Koike, Phys. Rev. D 48 (1993) 2313.
- [4] C.A. Dominguez and M. Loewe, Z. Phys. C (Particles & Fields) 51 (1991) 69; *ibid.* 58 (1993) 273; Phys. Lett. B 481 (2000) 295.
- [5] C.A. Dominguez, M. Loewe and J.C. Rojas, J. High Energy Phys. 0708 (2007) 040; E.V. Veliev and G. Kaya, arXiv:0902.3443.
- [6] C.A. Dominguez and M. Loewe, Z. Phys. C (Particles & Fields) 49 (1991) 423.
- [7] C.A. Dominguez, M. Loewe and J.S. Rozowsky, Phys. Lett. B 335 (1994) 506; C.A. Dominguez, M. S. Fetea and M. Loewe, Phys. Lett. B 387 (1996) 151; *ibid* B 406 (1997) 149; C.A. Dominguez, M. Loewe and C. van Gend, Phys. Lett. B 429 (1998) 64; *ibid* B 460 (1999) 442.
- [8] For recent results see e.g. H. Ohno *et al.*, arXiv:08103066 and references therein.
- [9] T. Altherr and D. Seibert, Phys. Rev. C 49 (1994) 1684.

- [10] P. Gerber and H. Leutwyler, Nucl. Phys. B 321 (1989) 387.
- [11] G. Boyd and D. E. Miller, arXiv:hep-ph/9608482 (unpublished); D.E. Miller, arXiv:hep-ph/0008031 (unpublished)
- [12] V.L. Eletsky, Phys. Lett. B 352 (1995) 440.
- [13] F. Klingl, S. Kim, S.H. Lee, P. Morath, and W. Weise, Phys. Rev. Lett. 82, 3396 (1999).
- [14] K. Morita and S.H. Lee, Phys. Rev. Lett. 100 (2008) 022301; arXiv:0711.3998.
- [15] Particle Data Group, C. Amsler *et al.*, Phys. Lett. B 667, 1 (2008).
- [16] R.A. Bertlmann, *et al.*, Z. Phys. C (Particles & Fields) 39 (1988) 231; C.A. Dominguez and J. Sola, Z. Phys. C (Particles & Fields) 40 (1988) 63; C.A. Dominguez and K. Schilcher, J. High Energy Phys. 01 (2007) 093.